



Residue® Complete

Guaranteed Analysis

Active Ingredient(s): as soil amending ingredient(s)

Microbial Content	0.62 %
<i>Bacillus amyloliquefaciens</i>	6.25×10^7 cfu/g
<i>Bacillus licheniformis</i>	2.50×10^8 cfu/g
<i>Bacillus megaterium</i>	6.25×10^7 cfu/g
<i>Bacillus pumilus</i>	1.88×10^9 cfu/g
<i>Phanerochaete chrysosporium</i>	4.38×10^4 cfu/g
<i>Trichoderma harzianum</i>	1.88×10^4 cfu/g
Sugars (Sucrose as microbial food)	74.3 %
Humic Acids (derived from Leonardite).....	5.0 %
Total Other Ingredients.....	20 %

Product Description:

Residue Complete is a proprietary blend of naturally occurring, nutrient cycling bacterial and fungal microorganisms plus a food source for digesting residual organic matter in or on the soil. These microorganisms were specially chosen to help aid in the breakdown of thatch and other organic matter in turf and reduce buildup of black layer on golf greens. It also contains soil conditioning humates to aid in improving soil conditions which enhance root vigor.

DPH Biologicals

21417 County Road 1950 East
Princeton, Illinois 61356 USA
Phone: (800) 648-7626
www.dphbiologicals.com

Net Weight: 30 lb. (13.63 kg)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

P-3-LBL-RC30TF-001

FIRST AID

Eye Contact: Immediately flush plenty of water. After initial flushing, remove any contact lenses and continue flushing for at least 15 minutes. If eye irritation persists: Get medical advice/attention. **Skin Contact:** Wash off immediately with plenty of soap and plenty of water while removing all contaminated clothes and shoes. If skin irritation persists, call a physician. **Inhalation:** Remove exposed individual(s) to fresh air for 20 minutes. Consult a physician/poison center if individuals condition declines or if symptoms persist. **Ingestion:** Rinse mouth. Do NOT induce vomiting. Drink 1 or 2 glasses of water. Never give anything by mouth to an unconscious person. Call a poison center or doctor/physician if you feel unwell.

Storage & Disposal:

Storage: Residue Complete should be stored in its original container, away from heat and direct sunlight. Preferably store in cool areas out of direct sunlight, away from children and pets, feed, and food products. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Disposal: Do not reuse this container. Dispose of contents/container in accordance with local/regional/national regulations.

Expiration Date: product guaranteed effective up to 2 years after date stamped on container.

Condition of Sale and Warranty: DPH Biologicals warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. Handling, storage and use of the product by Buyer or User are beyond the control of DPH Biologicals and Seller. Risks such as crop injury or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, or failure to follow label directions will be assumed by Buyer or User. IN NO CASE WILL DPH BIOLOGICALS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.



P-F-PWD-RCOMP-30LB

Mixing and Application:

Tank mix Residue Complete according to the directions in the table below. Mix only enough for the immediate application. Do not mix product with bactericides or copper compounds.

Liquid Solutions: it is recommended to mix Residue Complete with other products the day of use and to apply the product within 24 hours of wetting. It is recommended to utilize a 50-mesh screen in your spray unit to prevent clogging of your system. Consult your sales representative for more specific rate recommendations.

Dry Blending: blend Residue Complete with the appropriate amount of dry fertilizer or solid material to cover the acreage you intend to treat.

TURF APPLICATIONS		
Application	Rate	Frequency/Directions
Regular Maintenance	Apply 12 – 16 oz of product into 50 – 100 gallons of water per acre. Metric: Apply 340 – 454 g of product into 467 – 934 L of water per ha.	Apply monthly throughout growing season.
Overseeding, Newly Seeded Areas, Sod, Aeration	Apply 1.5 – 2 oz of product per 5,000 ft ² . Metric: Apply 43 – 57 g of product per 464 m ² .	Apply at time of aeration, sodding or seeding in the seed hopper or with starter fertilizer and reapply monthly to maturity.
Hydroseeding	Apply 1.5 – 2 oz of product per 5000 – 8000 ft ² per 500 – 800 gal of slurry. Metric: Apply 43 – 57 g of product per 464 – 743 m ² per 1900 – 3000 L of slurry	Add to tank before seed and mulch and reapply monthly until mature.
Sand Topdressing or Divot & Ball Mark Repair	Apply 12 – 16 oz of product per cubic yard.	Mix thoroughly with sand just prior to use. Hand Repair: Lightly sprinkle into divot hole prior to replacement of turf.

GREENHOUSE, NURSERY & LANDSCAPE APPLICATIONS		
Application	Rate	Frequency/Directions
Ornamental Tree and Shrub Care	<u>Initial Application:</u> Apply 12 – 16 oz of product per 100 gal of water. Metric: Apply 340 – 454 g of product per 378 L of water. <u>Maintenance:</u> Apply 12 oz per 100 gal of water. Metric: Apply 340 g of product per 378 L of water.	Soil inject, drench or spray at appropriate rates to ensure proper soil penetration.
New Field Plantings and Landscape Transplants	Mix 12 – 16 oz of product per 100 gallons of water. Metric: Mix 340 – 454 g of product per 378 L of water	Drench or sprench at appropriate rates to ensure proper soil penetration.
Greenhouse Plants, Injection Systems	<u>Initial planting:</u> Mix in 1.5 – 2 oz of product per gal stock tank at 1:100. Metric: Mix 43 – 57 g of product per 4 L stock tank at 1:100 <u>Constant Feed:</u> Mix 2 oz of product per gal stock tank at 1:100 Metric: Mix 57 g of product per 4 L stock tank at 1:100	Drench or sprench at appropriate rates to ensure proper soil penetration. Repeat twice a month in high porosity bark mixes and once a month in low porosity peat mixes.
Annual and Perennial Flower Beds	Mix 12 – 16 oz of product per 100 gal of water. Metric: 340 – 454 grams of product per 378 L of water.	Drench or sprench at appropriate rates to ensure proper soil penetration. Repeat monthly in low CEC soils and every other month in mid to high CEC soils.
Granular Fertilizer Addition	Mix 2.5 – 10 oz of product per 2000 lb. of fertilizer Mix 78 – 313 g of product per 1000 kg of fertilizer.	Mix last into blender. Caution: do not allow dust control measures to remove product during blending.